

Swift follow-up of the gigantic TeV outburst of PKS 2155 – 304 in 2006

L. Foschini^{*}, G. Ghisellini[†], F. Tavecchio[†], A. Treves^{**}, L. Maraschi[†], M. Gliozzi[‡],
C.M. Raiteri[§], M. Villata[§], E. Pian[¶], G. Tagliaferri[†], G. Tosti^{||}, R.M. Sambruna^{††}, G.
Malaguti^{*}, G. Di Cocco^{*} and P. Giommi^{‡‡}

^{*}INAF/IASF-Bologna, Via Gobetti 101, 40129, Bologna (Italy)

[†]INAF/Osservatorio Astronomico di Brera, Via Brera 28, 20121 Milano (Italy)

^{**}Dipartimento di Scienze, Università dell'Insubria, Via Valleggio 11, 22100 Como (Italy)

[‡]George Mason University, Department of Physics and Astronomy, School of Computational Sciences, Mail
Stop 3F3, 4400 University Drive, Fairfax, VA 22030, USA

[§]INAF/Osservatorio Astronomico di Torino, Via Osservatorio 20, 10025 Pino Torinese (Italy)

[¶]INAF/Osservatorio Astronomico di Trieste, Via G.B. Tiepolo 11, 34131 Trieste (Italy)

^{||}Osservatorio Astronomico, Università di Perugia, Via B. Bonfigli, 06126 Perugia (Italy)

^{††}NASA/Goddard Space Flight Center, Code 661, Greenbelt, MD 20771, USA

^{‡‡}ASI Science Data Centre, Via G. Galilei, 00044 Frascati, (Italy)

Abstract. At the end of July 2006, the blazar PKS 2155 – 304 ($z = 0.116$) underwent a strong outburst observed at TeV energies by HESS (up to 17 Crab flux level at $E > 200$ GeV). The *Swift* satellite followed the evolution of the source for about one month. The data analysis – reported in another paper – has shown that, despite the violent activity at TeV energies, the synchrotron energy distribution increased in normalization, but only with a small shift in frequency. In the present work, we fit the broad-band spectrum with a log-parabolic model, to search for indications of intrinsic curvature, which in turn is usually interpreted as a signature of energy-dependent acceleration mechanisms of electrons.

Keywords: BL Lacertae objects: general – BL Lacertae objects: individual (PKS 2155-304)

PACS: 95.85.-e; 95.85.Nv; 98.54.Cm; 98.70.Qy

INTRODUCTION

The HESS Cerenkov Telescope alerted the astronomical community on July 27, 2006 about an increase of activity of the blazar PKS 2155 – 304 ($z=0.116$) in the TeV energy range [1]. The same night the blazar displayed flares up to 17 Crab ($E > 200$ GeV) on time scales of 5 minutes [2]. An immediate X-ray/UV/Optical follow-up with the *Swift* satellite was activated and the analysis of data is reported in [3]. A prompt increase of the X-ray flux without large spectral changes was revealed. Interestingly, the frequency of the synchrotron peak remained at values similar to those observed in the past, with low TeV activity [4]. Further details can be found in [3]. However, the event was so spectacular and anomalous that it deserves further attention.

SPECTRAL ANALYSIS WITH THE LOG-PARABOLIC MODEL

After an initial spectral fit of the X-ray data with power-law (PL) and broken power-law (BPL) models, we tried a log-parabolic model, which was already successfully applied to other blazars (e.g. [5], [6], [7]). This model is represented by the relationship:


$$F(E) = KE^{-a-b\log(E)} \quad (1)$$

where E is the energy, K the normalization, a is the spectral slope (photon index) at 1 keV, and b is the curvature. We applied this model to the *Swift* observations with more than 100 s of exposure on the X-ray Telescope (XRT), which are listed in Table 1 of [3]. The results of the fits are reported in Table 1, while the values for a and b versus the normalization K are shown in Fig. 1. The fits with the log-parabolic model are generally comparable to the values

TABLE 1. Summary for fits with *Swift*/XRT data

Date	a	b	F*	$\tilde{\chi}^2/\text{dof}$
Apr 16	2.4 ± 0.1	0.4 ± 0.3	0.85	1.28/40
Apr 26	2.4 ± 0.1	< 0.61	1.27	1.03/19
Jul 29/31	2.48 ± 0.01	0.44 ± 0.04	3.40	1.16/320
Aug 1	2.63 ± 0.06	0.48 ± 0.18	2.70	0.95/92
Aug 2	2.61 ± 0.02	0.42 ± 0.07	2.44	1.00/196
Aug 3	2.46 ± 0.02	0.45 ± 0.07	2.83	1.32/206
Aug 5	2.69 ± 0.06	0.43 ± 0.19	1.89	0.80/92
Aug 6	2.64 ± 0.09	< 0.29	1.68	0.93/51
Aug 8	2.64 ± 0.06	0.44 ± 0.19	2.14	0.75/90
Aug 10	2.58 ± 0.07	0.24 ± 0.22	2.08	0.87/69
Aug 12	2.8 ± 0.2	< 0.63	1.27	1.18/16
Aug 20	2.4 ± 0.2	< 0.65	1.14	1.14/24
Aug 22	2.8 ± 0.2	0.89 ± 0.62	1.42	1.20/24

* Observed flux in the 0.3 – 10 keV energy band [10^{-10} erg/cm²/s].

**FIGURE 1.** Curvature (*b*) and photon index (or spectral slope *a*) versus model normalization (*K*) of the log-parabolic model of Eq. (1) applied to *Swift*/XRT data in the 0.3 – 10 keV energy range.

obtained by using single or broken power law models reported in [3]. In a few cases the log-parabolic model gives a better fit than the PL and BPL. In other cases, the resulting curvature is consistent with 0, i.e. the log-parabola mimics a PL. However, the limited photon statistic does not allow us to draw firm conclusions.

In Fig. 1 we display the curvature (*top panel*) and the photon index (*bottom panel*) plotted versus the normalization. The constant trend revealed by the data confirms the findings reported in [3]: there was no significant spectral changes in the X-ray emission from PKS 2155 – 304 during the outburst (corresponding to increasing X-ray fluxes in Fig. 1).

REFERENCES

1. Benbow W., Costamante L., Giebels B. on behalf of the HESS Collaboration, 2006, *Atel* **867**
2. Aharonian F., Akhperjanian A.G., Bazer-Bachi A.R. et al., *ApJ Letters*, submitted
3. Foschini L., Ghisellini G., Tavecchio F., et al., *ApJ* **657**, L81–L83 (2007).
4. Chiappetti L., Maraschi L., Tavecchio F., et al., *ApJ* **521**, 552–560 (1999).
5. Giommi P., Capalbi M., Fiacchi M., et al., “A Catalog of 157 X-ray Spectra and 84 Spectral Energy Distributions of Blazars observed with BeppoSAX” in *Proceedings of Blazar Astrophysics with BeppoSAX and other Observatories*, edited by P. Giommi, E. Massaro and G.G.C. Palumbo, ASI SDC and ESA-ESRIN, Frascati, 2001, p. 63 [astro-ph/0209596].
6. Massaro E., Tramacere A., Perri M., et al., *A&A* **448**, 861–871 (2006).
7. Tramacere A., Giommi P., Massaro E., et al., *A&A*, accepted for publication [astro-ph/0611276].